

PRVPATENT- OCH REGISTRERINGSVERKET
PatentavdelningenE 00/1423
REC'D 07 DEC 2000

WIPO

PCT

4

**Intyg
Certificate**

Härmed intygas att bifogade kopior överensstämmer med de handlingar som ursprungligen ingivits till Patent- och registreringsverket i nedannämnda ansökan.

This is to certify that the annexed is a true copy of the documents as originally filed with the Patent- and Registration Office in connection with the following patent application.

(71) Sökande AstaCarotene AB, Gustavsberg SE
Applicant (s)

(21) Patentansökningsnummer 9903619-6
Patent application number

(86) Ingivningsdatum 1999-10-07
Date of filing

Stockholm, 2000-11-30

För Patent- och registreringsverket
For the Patent- and Registration Office

A. Södervall
Anita Södervall

Avgift
Fee

**PRIORITY
DOCUMENT**
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

**PATENT- OCH
REGISTRERINGSVERKET
SWEDEN**

Postadress/Adress
Box 5055
S-102 42 STOCKHOLM

Telefon/Phone
+46 8 782 25 00
Vx 08-782 25 00

Telex
17978
PATOREG S

Telefax
+46 8 666 02 86
08-666 02 86

110013400/BN

Use and method of treatment

The present invention relates to the use and method of treatment concerning utilization of xanthophylls, e.g. astaxanthin, for induction of suppression of harmful Th1 mediated immune responses.

Background of the invention.

CD4 T lymphocytes can be subdivided into two major subsets - Th1 cells and Th2 cells. These cells release different sets of cytokines which define their distinct actions in immunity. Th1 cells secrete interferon-gamma (IFN- γ) and are mainly involved in activating macrophages and CD8+ cytotoxic T-lymphocytes. Th2 cells secrete the interleukins IL-4, IL-5 and IL-10 and are mainly involved in stimulating B cells to produce antibodies.

Immune modulation aims at altering the balance between different subsets of responding T cells so that damaging responses are suppressed. In many cases autoimmune diseases and intracellular infections are associated with the activation of Th1 cells, which activate macrophages and drive an inflammatory immune response. The drugs currently used to suppress the immune system can be divided into three categories:

1) Powerful anti-inflammatory drugs of the corticosteroid family such as prednisone. Glucocorticoids influence virtually every cellular and humoral mechanism related to inflammation and immune response. However, there are also many adverse effects, including fluid retention, weight gain, diabetes, bone mineral loss and thinning of the skin.

2) Cytotoxic drugs such as azathioprine and cyclophosphamide. Cytotoxic drugs cause immunosuppression by killing dividing cells and they have serious side-effects. The use of these compounds is limited due to a range of toxic effects on tissues which have continuous cell dividing, such as the bone marrow.

3) Cyclosporin A, tacromycin and rapamycin are powerful immunosuppressive agents that interfere with T-cell signaling.

All of these drugs are very broad in their action and inhibit protective functions of the immune system as well as pathological responses that cause tissue injury. Opportunistic infection is therefore a common complication of immune suppressive drugs.

It would be desirable to have an immunosuppressive agent that targets the specific part of the immune response that causes tissue injury. In particular, it would be desirable to obtain a medicament for induction of suppression of harmful Th1 mediated immune responses.

Description of the invention

The present invention provides a medicament for induction of suppression of harmful Th1 mediated immune responses.

One aspect of the invention is directed to the use of at least one type of xanthophylls for induction of suppression of harmful Th1 mediated immune responses.

It may be desirable in some cases to suppress harmful Th1 mediated immune responses also in patients who do not suffer from an infection or disease. For example, it would be beneficial to suppress the mediated immune responses before and after receiving a transplant to counteract rejection of the transplant or before and after receiving implantation of fertilized eggs in the uterus.

In a preferred embodiment of the invention the harmful Th1 mediated immune responses are caused by at least one disease from the group of autoimmune diseases and chronic viral and intracellular bacterial infections.

Examples of diseases that cause harmful mediated immune responses are Psoriasis vulgaris, Multiple sclerosis (MS), Rheumatoid arthritis, Crohn's disease, Insulin-dependant diabetes mellitus, Tuberculosis (TB), Acute graft-versus-host disease (transplant rejection) and HIV virus infection.

Xanthophylls, including astaxanthin, is a large group of carotenoids containing oxygen in the molecule in addition to carbon and hydrogen. The carotenoids are produced *de novo* by plants, fungi and some bacteria [Johnson E.A. and Schroeder W.A., 1995, Adv In Biochem Engin. Biotechn. 53: 119-178].

In a preferred embodiment of the invention, the type of xanthophyll is astaxanthin, preferably in a form esterified with fatty acids.

In a particularly preferred embodiment the astaxanthin is derived from a natural source, such as a culture of the algae *Haematococcus sp.*, e.g. *Haematococcus pluvialis*.

The medicament in the invention is preferably an oral preparation, which optionally comprises an oil of food grade and it is suitably presented in separate unit doses.

The medicament may comprise a mixture of different types of xanthophylls or different forms of the same xanthophyll, such as a mixture of synthetic astaxanthin and naturally produced astaxanthin.

The oral preparation may comprise in addition to the xanthophylls auxiliary ingredients which are pharmacologically acceptable inactive or active ingredients, such as flavoring agents, fillers, emulsifiers, etc.

Examples of separate unit doses are tablets, gelatin capsules and predetermined amounts of solutions, e. g. oil solutions, or emulsions, e.g. water-in- oil or oil-in-water emulsions.

Another aspect of the invention is directed to a method of inducing suppression
5 of harmful Th1 mediated immune responses in a human patient comprising administration of an response suppressing amount of at least one type of xanthophyll to said patient.

The examples and preferred embodiments described for the use aspect of the invention also apply for this method aspect of the invention.

In particular, harmful Th1 mediated immune responses are caused by at least
10 one disease from the group of autoimmune diseases and chronic viral and intracellular bacterial infections, such as Psoriasis vulgaris, Multiple sclerosis (MS), Rheumatoid arthritis, Crohn's disease, Insulin-dependent diabetes mellitus, Tuberculosis (TB), Acute graft-versus-host disease (transplant rejection) and HIV virus infection, and the type of xanthophyll is preferably astaxanthin, particularly in a form esterified with fatty acids, e.g. from a natural
15 source, such as a culture of the algae *Haematococcus sp.*

The daily doses of the active ingredient of the invention will normally be in the range of 0.01 to 10 mg per kg body weight for a human calculated on the amount of astaxanthin, but the actual dose will depend on the immune response of the individual human patient, the reason for suppression of the harmful Th1 mediated immune response, such as the
20 type of disease causing the enhanced pathological Th1 response, and the recommendations of the manufacturer.

The xanthophyll astaxanthin is commercially produced via culturing of the algae *Haematococcus sp.* by AstaCarotene AB, Gustavsberg, Sweden. It is marketed and sold in Sweden as a dietary supplement

25 Astaxanthin from other sources, and other xanthophylls as well, are expected to be similarly useful for the purposes of the invention. An advantage of using astaxanthin from algae is, however, that the astaxanthin exists in a form esterified with fatty acids [Renström B. et al, 1981, Phytochem 20(11) :2561-2564], which esterified astaxanthin thereby is more stable during handling and storage than free astaxanthin.

30 The naturally produced astaxanthin can be obtained also from fungi and crustaceans, in addition to from algae [Johnson E.A. and Schroeder W.A., *ibid*].

Case studies

During the last five years reports have been received from patients taking the commercial dietary supplement capsules of the algal meal of *Haematococcus pluvialis*, Astaxin®, containing 4 mg astaxanthin. The daily doses recommended as an antioxidant is one capsule per day. However, 2 – 6 times that dose has been used by some patients without adverse effects. On the contrary, the higher doses have been experienced as beneficial in alleviating symptoms associated with some chronic diseases.

Six patient histories are disclosed more in detail below.

Crohn's disease

10 Patient 1. Boy, 17 years old, who had suffered from Crohn's disease for at least four years. He has been treated with anti-inflammatory agents, such as cortisone. He started to take the commercial product Astaxin (two capsules, each containing 4 mg of astaxanthin, per day). In about two months the cortisone treatment was phased out and later on stopped altogether. The patient was asymptomatic for more than a year when he experienced a
15 relapse. He was then received a short-term treatment with cortisone in combination with Astaxin, and the cortisone treatment was again phased out.

20 Patient 2. Woman, about 50 years of age, who had suffered from Crohn's disease for a long time. She received treatment with cortisone. Now she has started to take Astaxin in parallel with her steroid medication and she reports that she feels considerably better.

25 Patient 3. Man, 48 years old, who has suffered from Crohn's disease for the last 20 years. He has been operated on several times and he has been treated with cortisone. Directly after the last operation he started taking Astaxin (6 capsules per day) and no cortisone. With regard to the circumstances, he has been asymptomatic. He has compared his clinical status after the operation with the status of two other patients who were operated on at the same time and who received conventional treatment with cortisone. In comparison with these two other patients his recovery has been fully equal with theirs, with the positive exception that edema in his colon diminished more quickly than in the two other patients.

Lichen ruber planus.

30 Patient 4. Woman, more than 70 years of age, who had suffered from the disease for several years. The symptoms of the disease were *inter alia* open wounds which had not healed. She had been treated with anti-inflammatory agents, such as cortisone, for several years, orally and also by injection directly to the local inflammation areas. The treatment has not led to any result. She started to take 4 capsules of Astaxin per day , and after some weeks

visible alleviation of the symptoms started to show up. The wounds were healed in slightly more than one month. During this period, the patient herself phased out the cortisone treatment. The dose of Astaxin was lowered to 2 capsules per day when she was asymptomatic. However, the symptoms returned in connection with a common cold. The dose was then
5 increased to 4 capsules per day and the wounds healed again. She says herself that she now feels considerably better.

Psoriasis.

Patient 5. Male, 40 years, who suffers from psoriasis and mainly shows itself in rough skin on the elbows. After treatment with a skin cream enriched with algal
10 meal/astaxanthin (100 mg astaxanthin /kg cream) twice a day for three weeks, the symptoms diminished.

Patient 6. Woman, 45 years old, who suffers from psoriasis and mainly shows itself in rough skin on the elbows. After treatment with a skin cream enriched with algal
15 meal/astaxanthin (100 mg astaxanthin /kg cream) twice a day for three weeks, the symptoms diminished.

Thus, positive reports have been received from several patients suffering from Crohn's disease, rheumatoid arthritis, psoriasis and lichen planus. All of these diseases are autoimmune diseases which are known to be Th1 cell mediated diseases.

Therefore it is likely that the Th1 mediated response in the patients has been
20 suppressed and that there is a shift of the Th1/Th2 balance of the immune response towards the Th2 response. Further, it is likely that patients suffering from other predominantly Th1 mediated diseases would benefit from suppression of the harmful Th1 response, as would patients undergoing transplantation or implantation of fertilized eggs in the uterus.

110013400/BN

Claims

1. Use of at least one type of xanthophylls for the production of a medicament for induction of suppression of harmful Th1 mediated immune responses.
- 5 2. Use according to claim 1, wherein the harmful Th1 mediated immune responses are caused by at least one disease from the group of autoimmune diseases and chronic viral and intracellular bacterial infections.
3. Use according to claim 2, wherein the disease is Psoriasis vulgaris, Multiple sclerosis (MS), Rheumatoid arthritis, Crohn's disease, Insulin-dependant diabetes mellitus, 10 Tuberculosis (TB), Acute graft-versus-host disease (transplant rejection), or HIV virus infection.
4. Use according to any one of claims 1 -3, wherein the type of xanthophyll is astaxanthin.
5. Use according to claim 4, wherein the astaxanthin is in a form esterified with 15 fatty acids.
6. Use according to claim 4 or 5, wherein the astaxanthin is derived from a natural source.
7. Use according to claim 6, wherein the natural source is a culture of the algae *Haematococcus sp.*
- 20 8. Use according to any one of the claims 1 - 7, wherein the medicament is an oral preparation.
9. A method of inducing suppression of harmful Th1 mediated immune responses in a human patient comprising administration of an Th1 response suppressing amount of at least one type of xanthophyll to said patient.
- 25 10. The method according to claim 9, wherein the harmful Th1 mediated immune responses are caused by at least one disease from the group of autoimmune diseases and chronic viral and intracellular bacterial infections.
11. The method according to claim 10, wherein the disease is Psoriasis vulgaris, Multiple sclerosis (MS), Rheumatoid arthritis, Crohn's disease, Insulin-dependant diabetes mellitus, Tuberculosis (TB), Acute graft-versus-host disease (transplant rejection), or HIV 30 virus infection.
12. The method according to claim 9, wherein the type of xanthophyll is astaxanthin.

13. The method according to claim 12, wherein the astaxanthin is in a form esterified with fatty acids.

14. The method according to claim 12 or 13, wherein the astaxanthin is derived from a natural source.

5 15. The method according to claim 14, wherein the natural source is a culture of the algae *Haematococcus sp.*

1
2
3
4
5
6
7
8
9
10

110013400/BN

Abstract

The use of at least one type of xanthophylls for the production of a medicament for induction of suppression of harmful Th1 mediated immune responses is disclosed.

- 5 Harmful Th1 mediated immune responses are caused by such autoimmune diseases and chronic viral and intracellular bacterial infections as Psoriasis vulgaris, Multiple sclerosis (MS), Reumatoid arthritis, Crohn's disease, Insulin-dependant diabetes mellitus, Tuberculosis (TB), Acute graft-versus-host disease (transplant rejection) and HIV virus infection. The preferred type of xanthophyll is astaxanthin, particularly in a form esterified with fatty acids,
- 10 obtainable by for example culturing the algae *Haematococcus sp.*

Further, a method of inducing suppression of harmful Th1 mediated immune responses in a human patient is disclosed.

